

# On Edge Dominating Number of Tensor Product of Cycle and Path

Robiatul Adawiyah<sup>1</sup>, Darmaji<sup>2</sup>, Reza Ambarwati<sup>1</sup>, Lela Nursafriada<sup>1</sup>, Inge Wiliandani Setya Putri<sup>1</sup>, Ermita Rizki Albirri<sup>1</sup>

<sup>1</sup>Department of Mathematics Education, University of Jember, Indonesia

<sup>2</sup>Department of Mathematics, Institut Teknologi Sepuluh Nopember, Indonesia

**Abstract**—A subset  $S'$  of  $E(G)$  is called an edge dominating set of  $G$  if every edge not in  $S'$  is adjacent to some edge in  $S'$ . The edge dominating number of  $G$ , denoted by  $\gamma'(G)$ , of  $G$  is the minimum cardinality taken over all edge dominating sets of  $G$ . Let  $G_1(V_1, E_1)$  and  $G_2(V_2, E_2)$  be two connected graph. The tensor product of  $G_1$  and  $G_2$ , denoted by  $G_1 \otimes G_2$  is a graph with the cardinality of vertex  $|V| = |V_1| \times |V_2|$  and two vertices  $(u_1, u_2)$  and  $(v_1, v_2)$  in  $V$  are adjacent in  $G_1 \otimes G_2$  if  $u_1 v_1 \in E_1$  and  $u_2 v_2 \in E_2$ . In this paper we study an edge dominating number in the tensor product of path and cycle. The results show that  $\gamma'(C_n \otimes P_2) = \lceil \frac{2n}{3} \rceil$  for  $n$  is odd,  $\gamma'(C_n \otimes P_3) = n$  for  $n$  is even, and the edge dominating number is undefined if  $n$  is odd. For  $n \in$  even number, we investigated the edge dominating number of its component on tensor product of cycle  $C_n$  and path. The results are  $\gamma'_c(C_n \otimes P_2) = \lceil \frac{n}{3} \rceil$  and  $\gamma'_c(C_n \otimes P_3) = \lceil \frac{n}{2} \rceil$  which  $C_n, P_2$  and  $P_3$ , respectively, is Cycle order  $n$ , Path order 2 and Path order 3.

**Keywords**—edge dominating number, tensor product, path, cycle.

## I. INTRODUCTION

One of the interesting topics in graph theory is dominating of graph. In recent years, there are some kinds of dominating in graph have been investigated. Most of those belong to the vertex dominating of graph. A few results have been obtained about the edge dominating of graph. In this paper, we mainly discuss about the edge dominating on some product operations of graph.

A subset  $H$  of  $E(G)$  is called an edge dominating set of  $G$  if every edge not in  $H$  is adjacent to some edge in  $H$ . The edge dominating number  $\gamma'(G)$  of  $G$  is the minimum cardinality taken over all edge dominating sets of  $G$ . The concept of edge dominating number are introduced by Mitchell and Hedetniemi [4] and was studied by several researches, such as V.R. Kulli [6,7,8,9,10], S. Arumugam and S. Velamal [1], Araya Chemchan [2], etc.

In graph theory, we have some graph operations, one of them is tensor product of graph. Let  $G_1(V_1, E_1)$  and  $G_2$

$(V_2, E_2)$  be two connected graph. The tensor product of  $G_1$  and  $G_2$ , denoted by  $G = G_1 \otimes G_2$  is a graph with the cardinality of vertex  $|V| = |V_1| \times |V_2|$  and two vertices  $(u_1, u_2)$  and  $(v_1, v_2)$  in  $V$  are adjacent in  $G_1 \otimes G_2$  if  $u_1 v_1 \in E_1$  and  $u_2 v_2 \in E_2$ . All graph considered here are finite, nontrivial, undirected, connected, without loops and multiple edges. We will verify the tensor product operation between cycle and path.

Cycle is a single vertex with a self-loop or simple graph  $C$  with  $|V_c| = |E_c|$  that can be drawn so that all of its vertices and edges lie on a single circle. A  $n$ -vertex cycle graph is denoted by  $C_n$  [3]. A path is a simple graph with  $|V_p| = |E_p| + 1$  that can be drawn so that all of its vertices and edge lie on a single straight line. A path graph with  $m$  vertices and  $m-1$  edges is denoted by  $P_m$  [3]. From the previous research, S. Arumugam and S. Velamal show that the edge dominating number of cycle graph is  $\gamma'(C_n) = n$  for  $n \geq 3$  [1] and based on the paper "The Neighborhood Total Edge dominating Number of A

Graph" by V.R. Kulli, we know that the edge dominating number of the path graph  $\gamma'(P_n) = n$  for  $n \geq 2, n \equiv 0 \pmod{3}$  [10]. In this paper, we are focus on finding the edge dominating number on tensor product of graph, respectively tensor product of cycle  $C_n$  and path which haven't been investigated before.

## II. EDGE DOMINATING NUMBER

An edge in graph  $G$  dominates itself and its adjacent. A set of edges  $S'$  in graph  $G$  is an edge dominating set, if each edge of  $G$  is dominated by some edges in  $S'$ . The dominating number  $\gamma'(G)$  of  $G$  is the minimum cardinality of edge dominating set of  $G$ . The concept of edge dominating was firstly introduced by Mitchell and Hedetniemi [4]. Let we take an example in cycle  $C_9$  (see figure 2), we can verify some edge dominating sets such as  $X_1 = \{e_1, e_3, e_4, e_6, e_8\}$ ,  $X_2 = \{e_1, e_4, e_6, e_8\}$  and  $X_3 = \{e_1, e_4, e_7\}$ . The cardinality of  $X_1, X_2$  and  $X_3$  are respectively 5, 4, and 3. Based on the definition, the dominating number of  $C_9$  is the minimum cardinality of edge dominating set of  $C_9$ . So, we get  $\gamma'(C_9) = 3$ .




Fig.1: cycle graph order 9

Consider the graph  $C_9$  shown in figure 1, we can see that the edge dominating set of  $C_9$  with cardinality 3 is not only  $X_3 = \{e_1, e_4, e_7\}$ . There also exist another edge dominating sets with cardinality 3, such that  $X_4 = \{e_2, e_5, e_8\}$  and  $X_5 = \{e_3, e_6, e_9\}$ . Our concern in this paper is not about the element of edge dominating set, but the minimum cardinality taking on edge dominating set. As long as the cardinality of edge dominating set is minimum, it can be concluded as edge dominating number of graph  $G$ . In order to verify the edge dominating number on tensor product of path and cycle in this paper, we need the following term. The degree of an edge  $e = uv$  of  $G$  is defined by  $deg(e) = deg(u) + deg(v) - 2$  [1]. For a real number  $x$ , the greatest integer less than or equal to  $x$  is denoted by  $\lfloor x \rfloor$  and the smallest integer greater than or equal to  $x$  is denoted by  $\lceil x \rceil$ .

### III. MAIN RESULTS

The following theorems show the properties of the edge dominating number on tensor product of path and cycle. Path order 2, path order 3, and cycle graph order  $n$ , respectively, are denoted by  $P_2$ ,  $P_3$ , and  $C_n$ . Here are some results of edge dominating number on tensor product of cycle and path.

**Theorem 3.1** Let  $C_n$  and  $P_2$  be, respectively cycle order  $n$  and path order 2. For  $n \geq 3$  and  $n$  is odd,

$$\gamma'(C_n \otimes P_2) = \lceil \frac{2n}{3} \rceil$$

**Proof.** The graph  $C_n \otimes P_2$  is a regular graph 2. By definition, for any regular graph with degree 2 is a cycle. If we observe graph  $C_n \otimes P_2$ , this graph is isomorphic with cycle graph with the cardinality of edges is  $2n$ . Thus, we should consider the edge dominating number of cycle graph. Because every edge in  $C_n \otimes P_2$  has degree 2, So, for every  $E \in S'$ ,  $E$  can dominated maximum 3 sides in  $C_n \otimes P_2$ ,  $|S'| \geq \frac{2n}{3}$ . Let says that the minimum cardinality of edge dominating number is  $\gamma'(C_n \otimes P_2) = \lceil \frac{2n}{3} \rceil$  for  $n \geq 3$ . In order to prove that  $\lceil \frac{2n}{3} \rceil$  is the minimum cardinality of the dominating set  $S'$  in  $C_n \otimes P_2$ , we use contradiction. Let  $S$  is

a minimum edge dominating set with cardinality  $\lfloor \frac{2n}{3} \rfloor - 1$ .

The maximum edge can be dominated is  $3(\lfloor \frac{2n}{3} \rfloor - 1) \leq 3(\frac{2n+2}{3}) - 1 = 2n - 1$ . We know that  $\|C_n \otimes P_2\| = 2n$ . If the cardinality of minimum edgedominating set  $S'$  is  $\lfloor \frac{2n}{3} \rfloor - 1$ , there are one edge which cannot be dominated by  $S'$ . So,  $|S'| = \gamma'(C_n \otimes P_2) = 2n$  is the minimum cardinality taken all of the dominating set. It can be concluded  $\gamma'(C_n \otimes P_2) = \lceil \frac{2n}{3} \rceil$ .

For  $n \geq 3$  and even, there is no domination number since  $(C_n \otimes P_2)$  is disconnected graph. We can find the edge dominating number only on connected graph. For  $n \in$  even number, the tensor product  $C_n \otimes P_2$  consist of two component of graph which is isomorphic one and other. If we separate the two component, we can see that each graph is isomorphic to cycle graph  $C_n$ . Then, we can verify a new theorem about edge dominating number on component of tensor product  $C_n \otimes P_2$  (denoted by  $\gamma'_c$ ), for  $n \geq 3$  and even.

**Theorem 3.2** Let  $C_n$  and  $P_2$  be, respectively, cycle order  $n$  and path order 2. For any component of  $C_n \otimes P_2$ , for  $n \geq 3$  and even,

$$\gamma'_c(C_n \otimes P_2) = \lceil \frac{n}{3} \rceil$$

**Proof.** We have noticed that the component of  $C_n \otimes P_2$  for  $n \in$  even number is isomorphic to cycle graph order  $n$ . Thus, we should consider the edge dominating number of cycle graph. Because every edges in cycle graph has degree 2, So, for every  $E \in S'$ ,  $E$  can dominated maximum 3 sides in  $C_n$ ,  $|S'| \geq \frac{n}{3}$ . Because  $n \in \mathbb{Z}^+$ , which  $\mathbb{Z}^+$  is denoted an integer more than zero, so  $S' = \lceil \frac{n}{3} \rceil$ .

Now, we proof  $\gamma'(C_n) \geq n$ . By contradiction, let  $|S'| = \gamma'(C_n) \leq \lfloor \frac{n}{3} \rfloor - 1$ . Without loose of generality, we can assume that  $S'$  is a minimum edge dominating set of cardinality  $\lfloor \frac{n}{3} \rfloor - 1$ . The maximum edge can be dominated is  $3(\lfloor \frac{n}{3} \rfloor - 1) \leq 3(\frac{n+2}{3}) - 1 = n - 1$ . Not all edges can be dominated by  $S'$ . So,  $|S'| = \gamma'(C_n) \leq \lfloor \frac{n}{3} \rfloor$  is the minimum cardinality taken all of the dominating set. We can conclude  $\gamma'_c(C_n \otimes P_2) = \lceil \frac{n}{3} \rceil$ , for any component of  $C_n \otimes P_2$  and  $n \in$  even number.

**Theorem 3.3** Let  $C_n$  and  $P_3$  be, respectively, cycle order  $n$  and path order 3. For  $n \geq 3$  and  $n$  is odd,

$$\gamma'(C_n \otimes P_3) = n$$

**Proof.** The tensor product of  $C_n \otimes P_3$  consist of  $2 \times n \times$

$(3 - 1) = 4n$  edges. The maximum edge degree in  $C_n \otimes P_3$  is  $deg(u_1v_1) = deg(u_1) + deg(v_1) - 2 = 4$ . So, every edge with maximum degree can dominate 5 other edges included itself. Suppose that, the edge with maximum degree is one of the element of edge dominating set in  $C_n \otimes P_3$ . There are  $\lfloor \frac{2n}{3} \rfloor$  edges with maximum degree which can dominate 5 edges included it's self. In order to determine the edge dominating number of  $C_n \otimes P_3$ , Let we observe the following condition of dominating set if the element of edge dominating set is the edge with maximum degree.

1. If  $n$  is multiple of three, there are  $\frac{2n}{3}$  edges which can dominate 2 edges and  $\lfloor \frac{2n}{3} \rfloor$  edges with maximum degree that can dominated 5 edges. The total edges can be dominated by those edges are  $5 \cdot \frac{2n}{3} + 2 \cdot \frac{2n}{3} = 4n$ . So, we need minimal  $|S'| = \frac{2}{3} + \frac{2n}{3} = n$  edges so that all edges in  $C_n \otimes P_3$  can be dominated. If the cardinality of  $S'$  is  $n - 1$ , there will be minimal 2 edges which can't be dominated by  $S'$ .

2. If  $n - 1$  is multiple of three, there are  $2 \left( \frac{n-1}{3} \right)$  edges which can dominate 2 edges,  $\lfloor \frac{2n}{3} \rfloor$  edges with maximum degree, and one edge that can dominate 4 other edges include it's self. So, the cardinality of edge dominating set is  $|S'| \geq \frac{2n-2}{3} + \frac{n-1}{3} + 1 = n$ . The total edges can be dominated by  $|S'| = n$  edges are  $5 \cdot \lfloor \frac{2n}{3} \rfloor + 2 \cdot \left( \frac{n-1}{3} \right) + 4 = 4n$ . If the cardinality of  $S'$  is  $n - 1$ , there will be minimal 2 edges which can't be dominated by  $S'$ .

3. If  $n + 1$  is multiple of three, there are  $2 \lfloor \frac{n}{3} \rfloor$  edges which can dominate 2 edges,  $\lfloor \frac{2n}{3} \rfloor$  edges with maximum degree, and one edge that can dominate 3 other edges include it's self. So, the cardinality of edge dominating set is  $|S'| \geq \frac{2n-1}{3} + \frac{n-2}{3} + 1 = n$ . The total edges can be dominated by  $|S'| = n$  edges are  $5 \cdot \lfloor \frac{2n}{3} \rfloor + 2 \cdot \left( \frac{n-1}{3} \right) + 4 = 4n$ . If the cardinality of  $S'$  is  $n - 1$ , there will be minimal 2 edges which can't be dominated by  $S'$ .

From those three conditions, we can see that the minimum cardinality of edge dominating set in each condition is  $n$ . So, it can be concluded  $\gamma'(C_n \otimes P_3) = n$ , for  $n \in \text{even number}$ .

For  $n \geq 3$  and even, there is no domination number since  $C_n \otimes P_3$  is disconnected graph. Because tensor product  $C_n \otimes P_3$  for  $n \in \text{even number}$  consist of disconnected graph, so the edge dominating number is undefined. Although we can not determine the edge dominating number of  $C_n \otimes P_3$ , but we can consider the component of  $C_n \otimes P_3$  to be analyzed. The tensor product

$C_n \otimes P_2$  consist of two component which is isomorphic. From the component of  $C_n \otimes P_3$ , we can determine the edge dominating number of each component in  $C_n \otimes P_3$ . Here is the theorem of edge dominating number on component of tensor product  $C_n \otimes P_2$  (denoted by  $\gamma'_c$ ), for  $n \geq 3$  and even.

**Theorem 3.4** Let  $C_n$  and  $P_3$  be, respectively, cycle order  $n$  and path order 3. For any component of  $C_n \otimes P_3$ , for  $n \geq 3$  and even,

$$\gamma'_s(C_n \otimes P_3) = \frac{n}{2}$$

**Proof.** The component of  $C_n \otimes P_3$  consist of  $2n$  edges. Suppose that, the edge with maximum degree is one of the element of edge dominating set in subgraph  $C_n \otimes P_3$ . The maximum edge degree in subgraph of  $C_n \otimes P_3$  is  $deg(u_1v_1) = deg(u_1) + deg(v_1) - 2 = 4$ . So, every edge with maximum degree can dominated 5 other edges included itself. If  $n$  is multiple of 3, there will be  $\frac{n}{3}$  edges with maximum degree that can dominate 5 edges and  $\frac{n}{6}$  which can dominate 2 edges. So, the total edge can be dominated by those edges are  $5 \cdot \frac{n}{3} + 2 \cdot \frac{n}{6} = 2n$ . We can say that  $\frac{n}{3} + \frac{n}{6} = \frac{n}{2}$  edges can dominate all edges in  $C_n \otimes P_3$ . If  $n - 1$  is multiple of 3, there will be  $\lfloor \frac{n}{3} \rfloor$  edges with maximum degree,  $\frac{n-4}{6}$  which can dominate 2 edges, and one edge which can dominated 3 edges. So, the total edge can be dominated by those edges are  $5 \cdot \lfloor \frac{n}{3} \rfloor + 2 \cdot \frac{n-4}{6} + 1.3 = 2n$ . There will be  $|S'| \geq \frac{n-1}{3} + \frac{n-4}{6} + 1 = \frac{n}{2}$  edges which can dominate all edges in  $C_n \otimes P_3$ . If  $n - 2$  is multiple of 3, there will be  $\lfloor \frac{n}{3} \rfloor$  edges which can dominate 5 edges,  $\frac{n-2}{6}$  which can dominate 2 edges, and one edge which can dominate 4 edges. So, the total edge can be dominated by those edges are  $5 \cdot \lfloor \frac{n}{3} \rfloor + 2 \cdot \frac{n-2}{6} + 1.4 = 2n$ . It means that  $|S'| \geq \frac{n-2}{3} + \frac{n-2}{6} + 1 = \frac{n}{2}$  edges can dominate all edges in  $C_n \otimes P_3$ . From those explanation, we can see that we need minimal  $\frac{n}{2}$  edges to dominate all edges in  $C_n \otimes P_3$ .

In order to check that  $|S'| = \frac{n}{2}$  is minimum cardinality of edge dominating set, we use a contradiction. Assume that  $|S'| = \frac{n}{2} - 1$  is minimum cardinality of edge dominating set. If  $|S'| = \frac{n}{2} - 1$ , there will be minimal two edges which can not be dominated. So, it can be concluded that  $\gamma'_s(C_n \otimes P_3) = \frac{n}{2}$ , for  $n \in \text{even number}$  and  $n \geq 3$ .

#### IV. CONCLUSION

In this paper, we have already investigated the edge dominating number on tensor product of cycle and path. The results are as follow:

- $\gamma'(C_n \otimes P_2) = \left\lceil \frac{2n}{3} \right\rceil$ , for  $n \geq 3$  and  $n$  is odd.
- $\gamma'_c(C_n \otimes P_2) = \left\lceil \frac{n}{3} \right\rceil$ , for any subgraph of  $C_n \otimes P_2$  and  $n \in \text{even number}$ .
- $\gamma'(C_n \otimes P_3) = 3$ , for  $n \geq 3$  and  $n$  is odd.
- $\gamma'_c(C_n \otimes P_3) = \frac{n}{2}$ , for any subgraph of  $C_n \otimes P_3$  and  $n \in \text{even number}$

#### REFERENCES

- [1] Arumugam,S., and Vellamal,S. (1998). Edge Domination In Graph, Taiwanese Journal of Mathematics vol.2(2) pp.173-179.
- [2] Chemcan, A. (2010). The Edge Domination Number of Connected Graphs. Australasian Journal of Combinatorics vol.48, 185-189.
- [3] Gross,L., Jonathan, Yellen, and Jay. (2006). Graph Theory and It's Application. United State Of America : Chapman and Hall CRC.
- [4] Hedetniemi,S. T., dan Mitchell, S. Edge Domination in Graphs. Congr Numer 19, 1977, pp. 489-509.
- [5] S.R.Jayaram. (1987) Line Domination in Graph. Graphs Combin vol 3 pp. 357-363.
- [6] V.R. Kulli. The Middle Edge Domination in Graph. J Computer and Mathematical Sciences vol.4, 2013, pp. 372-375.
- [7] V.R. Kulli. (2012). Inverse Total Edge Domination in Graph, In advances in Domination Theory I, V.R. Kulli ed. Vishwa Intenational Publication Gulbarga India pp. 35-44.
- [8] V.R. Kulli. (2012). The Edge Dominating Graph of a Graph, In advances in Domination Theory I, V.R. Kulli ed. Vishwa Intenational Publication Gulbarga India pp. 127-131.
- [9] V.R. Kulli.(2013)The Semientire Edge Dominating Graph. Ultra Scientist vol.25(3) A, 2013, pp. 431-434.
- [10] V.R. Kulli.(2015) The Neighborhood Total Edge Domination Number of A Graph. International Research Journal of Pure Algebra vol.5(3) A, pp 26-30.