

# Instruments for Diagnosing Stress in Nursing Professionals and Academics: A Systematic Review

Krislayne Veras Alexandre\*, Leonardo Moreira Rabelo\*\*, Gabriela Meira de Moura Rodrigues\*\*\*

\*Nursing Student - Midwest University Development Center, Luziânia-GO, Brazil.

Email: [krislaynev16@gmail.com](mailto:krislaynev16@gmail.com).

\*\*Nursing Student - Midwest University Development Center, Luziânia-GO, Brazil.

Email: [leomrstark@gmail.com](mailto:leomrstark@gmail.com).

\*\*\*Master in Biomedical Engineering - Midwest University Development Center, Luziânia-Go, Brazil.

Email: [professoragaby.meira@gmail.com](mailto:professoragaby.meira@gmail.com).

**Abstract— Introduction:** Stress is defined as feedback from the body in the face of psychophysiological stress. It can cause major health problems to the individual, directly affecting their quality of life in different ways, for there are some validated tools to detect symptoms and make their diagnosis. **Objective:** To point out instruments used to diagnose stress and to describe their differences. **Methodology:** Article conducted through systematic review, including studies that had greater relevance in their information and were related to health, and excluded those who escaped the proposed theme. In all, 32 sources were used, published between 1956 and 2019, thus covering initial stress research and more recent studies. **Results:** After searching the literature we found 8 validated instruments to diagnose stress, highlighting a translated version of 2017 of Stress Assessment in Nursing Students being the most current. **Discussions:** Stress is considered a disharmony situation of the organism, with adaptive responses that can be specific to the stressor and highly generalized. The importance of early diagnosis made by psychologists was highlighted, as well as the use of strategies to mitigate the effects of this disease. **Conclusion:** The means to diagnose stress are diverse, as stressors, so health professionals should be aware of all signs of stress experienced by both nursing students and trained nurses.

**Keywords**— Diagnosis, Stress, Nursing, Nursing Academics.

## I. INTRODUCTION

Stress is characterized as an organism's response to a threatening situation, that is, a psychophysiological exhaustion, which can cause major health problems and drastically affect an individual's quality of life in many ways. Studies show that 90% of the population is affected by this disease, and is then seen as a global epidemic that may eventually cause a rise in blood pressure [1][2][3].

In this context, nursing is classified as the fourth most stressful profession, precisely because it has a higher susceptibility to stressors, since this occupation directly witnesses the health / disease process, which often happens in precarious conditions, in addition to being read daily, with suffering and death [4][5].

The stress caused by occupational stress contributes to the development of another harm, Burnout syndrome, occupational exhaustion that acts as a response caused by the chronic stress that makes professionals sick, at times when combat strategies are poorly resolved, causing a deficit in the quality of care provided, as well as physical and psychological damage [6][5][7].

For some authors, overload, communication problems with staff, responsibility to provide adequate care, interpersonal conflicts and job dissatisfaction are also stressors present in nursing work [8]. In short, these stressors are closely linked to the inherent factors of work [9].

In universities, nursing students are constantly facing stressful situations, precisely because they carry

responsibilities that concern the lives and health of others. Given this it is noticeable that stress is rooted in the profession, both trained professionals and academics, due to the demand for large collections lived [10].

During nursing undergraduate stress is triggered when the student feels overwhelmed with academic demands, the beginning of contact with human pain and suffering in the internship field, insecurity about the professional future, in addition to the financial factor, since part of the students work to fund their studies. Symptoms affect their interpersonal relationships that compromise their well-being, so reducing these factors is necessary, since the labor market has received professionals with stress symptoms, which can interfere with the quality of nursing care offered to patient in the early years of formation [11].

This way, further studies are needed to provide a better understanding of stress diagnoses in order to develop possible prophylaxis methods for stress mitigation. Therefore, the objectives of this study are to point out the instruments used to diagnose stress, besides describing their differences.

## II. METHODOLOGY

*Chart 1 - Instruments to diagnose stress*

AUTHORS	INSTRUMENT	INSTRUMENT SUMMARY
[13].	Bianchi Stress Scale(annexl).	<ul style="list-style-type: none"><li>• Focused on Determining the Level of Stress in Hospital Nursing in the Basic Exercise of Her Work;</li><li>• It is self-applicable;</li><li>• It takes an average of 15 minutes to answer;</li><li>• Formed by 51 items, divided into 6 domains, which gets a score ranging from 1 to 7;</li><li>• DOMAINS:<ul style="list-style-type: none"><li>1 - relationship with other units and supervisors (nine items: 40, 41, 42, 43, 44, 45, 46, 50, 51);</li><li>2 - activities related to the proper functioning of the unit (six items: 1, 2, 3, 4, 5, 6);</li><li>3 - personnel management activities (six items: 7, 8, 9, 12, 13, 14);</li><li>4 - nursing care provided to the patient (fifteen items: 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30);</li><li>5 - coordination of the unit's activities (eight items: 10, 11, 15, 31, 32, 38, 39, 47);</li><li>6 - working conditions for the performance of nurses' activities (seven items: 33, 34, 35, 36, 37, 48, 49).</li></ul></li><li>• SCORES:<ul style="list-style-type: none"><li>≤ 3.0: low stress level;</li><li>Between 3.1 and 5.9: medium stress level;</li></ul></li></ul>

This article is based on a systematic review on Google Scholar, PubMed and Scielo platforms. This type of research is characterized by conducting a synthesis on a particular health theme, aiming to provide data obtained on a problem [12].

For the execution of this work, a wide search in the literature was performed. In the end 32 sources were used, published from 1956 to 2019, thus covering initial research on stress, as well as more recent studies. We selected scientific articles that presented greater relevance in their data and were related to the health area, and excluded those who escaped from the theme addressed. The instruments were left in their original languages to retain their original quirks and characteristics.

## III. RESULTS

Considering the focus and general objective of this study, we searched the world literature for instruments that can measure and diagnose stress. The data obtained were distributed in a table for better organization, and there is also a short synthesis about it, aiming to help researchers who wish to use any of these tools.

		≥ 6.0: high stress level.
[14].	Stress Assessment in Nursing Students(annex 2).	<ul style="list-style-type: none"> <li>• Aimed at nursing academics, but there is guidance for use on different samples;</li> <li>• The instrument consists of 30 items, organized into six domains: <ul style="list-style-type: none"> <li>1- Realizationofpracticalactivities;</li> <li>2- Professional communication;</li> <li>3- Time management;</li> <li>4- Environment;</li> <li>5- Professional qualification;</li> <li>6- Theoreticalactivity.</li> </ul> </li> </ul>
[14] <i>apud</i> [15].	Stress Assessment in Nursing Students – VersionReduced (annex 3).	<ul style="list-style-type: none"> <li>• Aimed at university nursing students;</li> <li>• It can be applied anywhere in the undergraduate degree;</li> <li>• This version consists of 19 items;</li> <li>• Consisting of 4 domains: <ul style="list-style-type: none"> <li>1- Realization of practical activities (Items 2, 3, 4 and 5);</li> <li>2- Theoreticalactivity (Items 1, 7, 9, 10, 15, 17 and 19);</li> <li>3- Environment (Items 8, 14, 16 and 18);</li> <li>4- Vocational Training (Items 6, 11, 12 and 13).</li> </ul> </li> </ul>
[16].	<i>Perceived Stress Scale</i> (PSS) (figures 1 and 2).	<ul style="list-style-type: none"> <li>• The scale has 14 questions, the answers are: 0: <i>never</i>; 1: <i>almost never</i>; 2: <i>sometimes</i>; 3: <i>fairly often</i>; 4: <i>very often</i>;</li> <li>• The items declared positive are: 4, 5, 6, 7, 9, 10 and 13.</li> </ul>
[16] <i>apud</i> [17].	Perceived Stress Scale in Brazilian Portuguese (annex 4).	<ul style="list-style-type: none"> <li>• The article aims to translate the Perceived Stress Scale (PSS) into Portuguese and measure the perceived stress in the elderly in Brazil;</li> <li>• The scale has 14 questions with answer options that can be: 0: <i>never</i>; 1: <i>almost never</i>, 2: <i>sometimes</i>; 3: <i>almost always</i>; 4: <i>always</i>;</li> <li>• Positive questions are: 4, 5, 6, 7, 9, 10 and 13.</li> </ul>
[16] <i>apud</i> [18].	Perceived Stress Scale in Portuguese from Portugal (annex 5).	<ul style="list-style-type: none"> <li>• PSS-10 includes items 1, 2, 3, 6, 7, 8, 9, 10, 11, and 14 of PSS-14;</li> <li>• Possible answers are: 0- <i>never</i>, 1- <i>almost never</i>, 2- <i>sometimes</i>, 3- <i>often</i>, and 4- <i>very often</i>).</li> </ul>
[16] <i>apud</i> [19].	<i>Perceived Stress Scale 4</i> (PSS-4) (annex 6).	<ul style="list-style-type: none"> <li>• PSS-4 is formed by the numbers 2, 6, 7 and 14 of PSS-14;</li> <li>• This scale can be used in telephone interviews and in other situations where a smaller scale is required to provide data under stress.</li> </ul>
[20] <i>apud</i> [1].	Lipp Adult Stress Symptom Inventory (annex7).	<ul style="list-style-type: none"> <li>• It consists of 53 symptoms, organized hierarchically by symptom intensity. It takes about 10 minutes to perform, and can only be done in one interviewee or in groups of up to 20 people. To respond to each of the phases, the person indicates the physical or psychological signals felt in the last 24 hours, the past week or the last month [21] [22].</li> <li>• This instrument contains 37 somatic and 19 psychological symptoms, which are often repeated, modifying only their intensity and severity;</li> <li>• It consists of three frames relating to the four phases of stress.  1st chart: has 15 items related to the alert phase, of which 12 relate to physical symptoms and 3 to psychological symptoms experienced by the person in the last 24 hours;  2nd chart: it consists of 15 items, related to the phases of resistance and near exhaustion, 10 physical and 5 psychological symptoms observed in the last week;  3rd chart: includes 23 items, which are equivalent to the exhaustion phase, of</li> </ul>

these 12 are physical and 11 psychological signs, experienced in the last month.

Source: Prepared by the authors.

After investigation of the literature, a total of 8 validated instruments were found, being the 2017 smallest version of Stress Assessment in Nursing Students the most current, already the oldest was PSS-14, 10 and 4, published in 1983 [16]. This scale was translated into Portuguese from Brazil [17] and Portugal, but in this study, we used one of the shortened versions, the PSS-10 [18].

Psychological assessments are systematic observational procedures that record the conduct and responses of individuals for the purpose of describing and / or measuring psychological peculiarities and processes, commonly understood in the areas of emotion / affect, cognition / intelligence, motivation, personality, psychomotor, attention, memory, perception, among others, in their various ways of expression, based on patterns defined by the construction of instruments [23].

Thus, it should be emphasized that the use of instruments for the assessment or measurement of psychological characteristics for the purpose of psychological diagnosis is a private matter of the psychologist [24].

#### **IV. DISCUSSIONS**

Originally from physics, the word stress was used in the early twentieth century by Hans Selye to name the joint actions of forces that can reach any part of the human body. He presented the thesis that a series of stereotyped psychological and physiological occurrences in critically ill patients reproduced serious and lasting consequences of adaptive responses. He referred to this as the 'General Adaptation Syndrome (GAS)' or 'Stress Syndrome (SS)' [25].

Stress can still be classified as a disharmony situation when the body's homeostasis is endangered, with adaptive responses that may or may not be stressor-specific [26].

If this problem is not taken care of, it affects the health of the individual, the efficiency and professional and personal performance and can cause a scenario of depression, but it is not simple to find the agent responsible for stress, so it is suggested that the best way to detect it is through diagnosis and consultation with able psychologists using appropriate instruments [27].

In a survey of 136 nursing students from a private institution, 30.9% showed low perception of stress, 46.3%

medium and 22.8% high. From these results we can evidence something positive related to the small number of students with high perception, because the available literature shows that in most cases the rates are higher [28].

As in a study that showed that among 209 academics, 163 had stress indicators and only 22% of the total did not show the indicative [29]. Authors say that the high perception of stress affects performance, lifestyle and causes damage to the health of the university, point out that the higher the perception of stress the greater the danger of illness [28].

From the relationship between stress in future nurses and sociodemographic factors, it was found that 75% of a sample of 32 people were classified with signs of this disease and the sociodemographic variable that showed the highest relationship with stress was marital status [30].

Because of the stressors caused by the course, academics use methods to relax and combat stress, but still use smoking and alcohol as a means to remedy their effects [31].

Coping is cited as a strategy to alleviate stress in nursing students. It is characterized by the ability to cope with and adapt to stressful events in a positive way, concerns the form of behavior or even thoughts in the face of such events, and can also be considered a union of physical and psychological techniques that can be learned, changed, and even discarded [10].

In nurses who perform the function, authors suggest that these are the most susceptible to stress at work, given that the deficit of professionals in the area causes an increase and overload of service, along with the separation relationship between nurses and technicians or nursing assistants, decreased professional authority and autonomy, and disagreements with other health officials [32].

#### **V. CONCLUSION**

As noted, the tools to diagnose stress are numerous, which makes a vast possibility for researchers. In addition, they can be applied to both active nursing professionals and academics, which increases the possibilities for future studies.

The stressors are varied, with different nature and several triggering factors. They are closely linked with the performance of nursing professionals, since the academic life, because students go through situations that can trigger imbalance due to both professional and social pressure, so

health professionals should pay attention to all symptoms of stress experienced by both academics and working nurses.


## REFERENCES

- [1] Gato RC. Estresse e sintomas em profissionais de enfermagem de um hospital universitário [monografia]. Manaus: Universidade Federal do Amazonas; 2014.
- [2] Hamer M. Malan L. Psychophysiological risk markers of cardiovascular disease. *Neurosci Biobehav Rev*. 2010;35(1):76-83.
- [3] Chandola T. Britton A. Brunner E. Hemingway H. Malik M. Kumari M. et al. Work stress and coronary heart disease: what are the mechanisms? *European Heart Journal*. 2008 mar;29(5):640-48.
- [4] Fonseca JRF. Estresse ocupacional em enfermeiros de unidades de emergência no município de Manaus/AM [dissertação]. Manaus: Universidade Federal do Amazonas; 2012.
- [5] Freitas EO. Estresse, Coping, Burnout, sintomas depressivos e Hardiness entre discentes de enfermagem [dissertação]. Santa Maria: Universidade Federal de Santa Maria; 2012.
- [6] Santana LL. Miranda FMDA. Karino ME. Baptista PCP. Felli VEA. Sarquis LMM. Cargas e desgaste de Trabalho vivenciados entre trabalhadores de saúde em um hospital de ensino. *Revista Gaúcha de Enfermagem*. 2013 mar;34(1):64-70.
- [7] Silva AA. Rotenberg L. Fischer FM. Nursing work hours: individual needs versus working conditions. *Revista de Saúde Pública*. 2011 dez;45(6):1117-26.
- [8] Miquelim JDL. Carvalho CBO. Gir E. Pelá NTR. Estresse nos profissionais de enfermagem que atuam em uma unidade de pacientes portadores de HIV-Aids. DST – J bras Doenças Sex Transm. 2004;16(3): 24-31.
- [9] Paschoalini B. Oliveira MM. Frigério MC. Dias ALRPD. Santos FH. Efeitos cognitivos e emocionais do estresse ocupacional em profissionais de enfermagem. *Acta Paul. Enferm*. 2008;21(3):487-92.
- [10] Hirsch CD. Barlem ELD. Almeida LK. BarlemJGT. FigueiraAB. Lunardi VL. Estratégias de coping de acadêmicos de enfermagem diante do estresse universitário. *Ver. Bras. Enferm*. 2015 set/out;68(5):783-90.
- [11] Santos RJLL. Sousa EP. Rodrigues GMM, Quaresma PC. Estresse em acadêmicos de enfermagem: importância de identificar o agente estressor. *Braz. J. Hea. Rev*. 2019 mar/abr;2(2):1086-94.
- [12] Ercole FF. Melo LS. Alcoforado CLGC. Integrative review versus systematic review. *Reme*. 2014 jan/mar;18(1):1-260.
- [13] Bianchi ERF. Escala Bianchi de Stress. *Revista da Escola de Enfermagem da Usp*. 2009 dez;43:1055-62.
- [14] Costa ALS. Polak C. Construção e validação de instrumento para Avaliação de Estresse em Estudantes de Enfermagem (AEEE). *Revista da Escola de Enfermagem da Usp*. 2009 dez;43:1017-26.
- [15] Costa ALS, Silva RM, Mussi FC, Serrano PM, Graziano ES, Batista KM. Short version of the “instrument for assessment of stress in nursing students” in the Brazilian reality. *Rev. Latino-Am. Enfermagem*. 2017;25.
- [16] Cohen S. Kamarck T. Mermelstein R. A global measure of perceived stress. *Journal of Health and Social Behavior*. 1983 dez;24(4):385-96.
- [17] LuftCDB. Sanches SO. MazoGZ. Andrade A. Versão brasileira da escala de estresse percebido: tradução e validação para idosos. *Rev. Saúde Pública*. 2007;41(4):606-15.
- [18] Trigo M. Canudo N. Branco F. Silva D. Estudo das propriedades psicométricas da Perceived Stress Scale (PSS) na população portuguesa. *Psychologica*. 2010;53:353-78.
- [19] The Ohio state university. Perceived stress scale 4 (PSS-4) [internet]. [201-?]. [acesso em 2019 maio 07] Disponível em: <http://ohnurses.org/wp-content/uploads/2015/05/Perceived-Stress-Scale-4.pdf>.
- [20] Lipp MEN. Guevara AJH. Validação empírica do inventário de sintomas de stress. *Estudos de Psicologia*. 1994 set;11(3):43-9.
- [21] Lipp MEN, Malagris LEN. Manejo do estresse. In B. Range (Org.), *Psicoterapia comportamental e cognitiva: pesquisa, prática, aplicação e problemas*. Campinas: Editorial Psy II; 2001.
- [22] Rossetti MO, Ehlers DM, Guntert IB, Leme IFAS, Rabelo ISA, Tosi SMVD, et al. O inventário de sintomas de stress para adultos de lipp (ISSL) em servidores da polícia federal de São Paulo. *Revista Brasileira de Terapias Cognitivas*. 2008; 4(2).
- [23] Brasil. Resolução CFP n.º 005/2012. Altera a Resolução CFP n.º 002/2003, que define e regulamenta o uso, a elaboração e a comercialização de testes psicológicos [internet]. Brasília; 2012. [acesso em 2019 abr 01]. Disponível em: <https://site.cfp.org.br/wp->

- content/uploads/2012/03/Resolucao\_CFP\_005\_12\_1.pdf.
- [24] Brasil. Presidência da República, Casa Civil, Subchefia para Assuntos Jurídicos. Lei nº 4.119, de 27 de agosto de 1962. Dispõe sobre os cursos de formação em psicologia e regulamenta a profissão de psicólogo [internet]. Brasília; 1962. [acesso em 2019abr01]. Disponível em: [http://www.planalto.gov.br/ccivil\\_03/leis/1950-1969/l4119.htm](http://www.planalto.gov.br/ccivil_03/leis/1950-1969/l4119.htm).
- [25] Selye H. The stress of life. New York: McGraw-Hill; 1956.
- [26] Chrousos G. Loriaux D. Gold P. Mechanisms of physical and emotional stress. New York: Plenum Press; 1988.
- [27] Martins MGT. Sintomas de Stress em Professores Brasileiros. Revista Lusófona de Educação. 2007;2(1):109-28.
- [28] Flauzino MM. Garcia VPT. Gonçalves LA. Correia C. Parmigiani RS. Pereira SS. et al. Percepção de estresse nos acadêmicos de enfermagem de uma instituição particular de ensino. In: 8º Sim Saúde-Simpósio em Saúde, 2017; Araçatuba. Faculdade de Odontologia de Araçatuba, UNESP; 2017.
- [29] Preto VA, Sailer GC, Pereira SS, Teixeira CC, Parmigiani RS. Investigação sobre estresse e fases do estresse em universitários de enfermagem. In: 8º Sim Saúde-Simpósio em Saúde, 2017; Araçatuba. Faculdade de Odontologia de Araçatuba, UNESP; 2017.
- [30] Preto VA. Benevides MS. Queiroz BG. Pereira SS. Souza BOP. Sailer GC. et al. Estresse e características sociodemográficas em universitários de enfermagem. Rev. enferm. UFPE on-line. 2018 mar;12(3):701-7.
- [31] Gouveia MTO. Santana HS. Santos AMR. Costa GR. Fernandes MA. Quality of life and welfare of nursing university students: integrative review. Rev. Enferm. UFPI. 2017 Jul/set;6(3):72-8.
- [32] Vagharseyyedin SA. Vanaki Z. Mohammadi E. The nature nursing quality of work life: an integrative review of literature. West J Nursing Res. 2011;33(6):786-804.

**Annex 1 - Bianchi Stress Scale [13].**

Este questionário tem a finalidade de levantar dados para conhecer a sua opinião quanto ao desempenho de suas atividades. NÃO PRECISA IDENTIFICAÇÃO. Assinale a alternativa que revele a sua percepção, levando em consideração os números:



*não se aplica ou com médio muito  
“não faço” desgastante desgastante*

**PARTE 1**

Sexo :                    feminino ( )                    masculino ( )

Faixa etária :            ( ) 20 a 30 anos  
                                  ( ) 31 a 40 anos  
                                  ( ) 41 a 50 anos  
                                  ( ) mais de 50 anos

Cargo:

Unidade a que pertence:

Tempo de formado:( ) menos de 1 ano  
                                  ( ) de 2 a 5 anos  
                                  ( ) de 6 a 10 anos  
                                  ( ) 11 a 15 anos  
                                  ( ) mais de 16 anos

Cursos de pós-graduação :( ) não  
                                  ( ) sim Qual (is)

Tempo de trabalho nessa unidade:

**PARTE 2**

1. Previsão de material a ser usado	0 1 2 3 4 5 6 7
2. Reposição de material	0 1 2 3 4 5 6 7
3. Controle de material usado	0 1 2 3 4 5 6 7
4. Controle de equipamento	0 1 2 3 4 5 6 7
5. Solicitação de revisão e consertos de equipamentos	0 1 2 3 4 5 6 7
6. Levantamento de quantidade de material existente na unidade	0 1 2 3 4 5 6 7
7. Controlar a equipe de enfermagem	0 1 2 3 4 5 6 7
8. Realizar a distribuição de funcionários	0 1 2 3 4 5 6 7
9. Supervisionar as atividades da equipe	0 1 2 3 4 5 6 7
10. Controlar a qualidade do cuidado	0 1 2 3 4 5 6 7
11. Coordenar as atividades da unidade	0 1 2 3 4 5 6 7
12. Realizar o treinamento	0 1 2 3 4 5 6 7
13. Avaliar o desempenho do funcionário	0 1 2 3 4 5 6 7
14. Elaborar escala mensal de funcionários	0 1 2 3 4 5 6 7
15. Elaborar relatório mensal da unidade	0 1 2 3 4 5 6 7
16. Admitir o paciente na unidade	0 1 2 3 4 5 6 7
17. Fazer exame físico do paciente	0 1 2 3 4 5 6 7

18. Prescrever cuidados de enfermagem	0 1 2 3 4 5 6 7
19. Avaliar as condições do paciente	0 1 2 3 4 5 6 7
20. Atender as necessidades do paciente	0 1 2 3 4 5 6 7
21. Atender as necessidades dos familiares	0 1 2 3 4 5 6 7
22. Orientar o paciente para o auto cuidado	0 1 2 3 4 5 6 7
23. Orientar os familiares para cuidar do paciente	0 1 2 3 4 5 6 7
24. Supervisionar o cuidado de enfermagem prestado	0 1 2 3 4 5 6 7
25. Orientar para a alta do paciente	0 1 2 3 4 5 6 7
26. Prestar os cuidados de enfermagem	0 1 2 3 4 5 6 7
27. Atender as emergências na unidade	0 1 2 3 4 5 6 7
28. Atender aos familiares de pacientes críticos	0 1 2 3 4 5 6 7
29. Enfrentar a morte do paciente	0 1 2 3 4 5 6 7
30. Orientar familiares de paciente crítico	0 1 2 3 4 5 6 7
31. Realizar discussão de caso com funcionários	0 1 2 3 4 5 6 7
32. Realizar discussão de caso com equipe multiprofissional	0 1 2 3 4 5 6 7
33. Participar de reuniões do Departamento de Enfermagem	0 1 2 3 4 5 6 7
34. Participar de comissões na instituição	0 1 2 3 4 5 6 7
35. Participar de eventos científicos	0 1 2 3 4 5 6 7
36. O ambiente físico da unidade	0 1 2 3 4 5 6 7
37. Nível de barulho na unidade	0 1 2 3 4 5 6 7
38. Elaborar rotinas, normas e procedimentos	0 1 2 3 4 5 6 7
39. Atualizar rotinas, normas e procedimentos	0 1 2 3 4 5 6 7
40. Relacionamento com outras unidades	0 1 2 3 4 5 6 7
41. Relacionamento com centro cirúrgico	0 1 2 3 4 5 6 7
42. Relacionamento com centro de material	0 1 2 3 4 5 6 7
43. Relacionamento com almoxarifado	0 1 2 3 4 5 6 7
44. Relacionamento com farmácia	0 1 2 3 4 5 6 7
45. Relacionamento com manutenção	0 1 2 3 4 5 6 7
46. Relacionamento com admissão/alta de paciente	0 1 2 3 4 5 6 7
47. Definição das funções do enfermeiro	0 1 2 3 4 5 6 7
48. Realizar atividades burocráticas	0 1 2 3 4 5 6 7
49. Realizar tarefas com tempo mínimo disponível	0 1 2 3 4 5 6 7
50. Comunicação com supervisores de enfermagem	0 1 2 3 4 5 6 7

#### Annex2–Stress Assessment in Nursing Students [14].

Leia atentamente cada item abaixo e marque com um “X” o número correspondente com a intensidade de estresse que a situação lhe provoca, conforme a legenda a seguir:

- 0 = não vivencio a situação
- 1 = não me sinto estressado com a situação
- 2 = me sinto pouco estressado com a situação
- 3 = me sinto muito estressado com a situação

Nova Numeração	Itens	0 1 2 3			
		0	1	2	3
1.	Ter preocupação com o futuro profissional				

2.	A obrigatoriedade em realizar os trabalhos extraclasse		
3.	Estar fora do convívio social traz sentimentos de solidão		
4.	Realizar os procedimentos assistenciais de modo geral		
5.	As novas situações que poderá vivenciar na prática clínica		
6.	Comunicação com os demais profissionais da unidade de estágio		
7.	O ambiente da unidade clínica de estágio		
8.	Comunicação com os profissionais de outros setores no local de estágio		
9.	Ter medo de cometer erros durante a assistência ao paciente		
10.	A forma adotada para avaliar o conteúdo teórico		
11.	Distância entre a faculdade e o local de moradia		
12.	Executar determinados procedimentos assistenciais		
13.	Sentir insegurança ou medo ao fazer as provas teóricas		
14.	O grau de dificuldade para a execução dos trabalhos extraclasse		
15.	A semelhança entre as situações que vivencia no estágio e aquelas que poderá vivenciar na vida profissional		
16.	Perceber as dificuldades que envolvem o relacionamento com outros profissionais da área		
17.	Pensar nas situações que poderá vivenciar quando for enfermeiro		
18.	Tempo reduzido para estar com os familiares		
19.	Perceber a responsabilidade profissional quando está atuando no campo de estágio		
20.	Observar atitudes conflitantes em outros profissionais		
21.	Sentir que adquiriu pouco conhecimento para fazer a prova prática		
22.	Transporte público utilizado para chegar à faculdade		
23.	Tempo exigido pelo professor para a entrega das atividades extraclasse		
24.	Distância entre a maioria dos campos de estágio e o local de moradia		
25.	Vivenciar as atividades, como enfermeiro em formação, no campo de estágio		
26.	Faltar tempo para o lazer		
27.	Perceber a relação entre o conhecimento teórico adquirido no curso e o futuro desempenho profissional		
28.	Assimilar o conteúdo teórico-prático oferecido em sala de aula		
29.	Transporte público utilizado para chegar ao local do estágio		
30.	Faltar tempo para momentos de descanso		

#### Cálculo dos escores:

Para aferição do resultado, deve ser feita a soma do número correspondente da intensidade de estresse dos itens presentes em cada domínio. O domínio com maior pontuação será considerado predominante e com maior intensidade de estresse para o respondente.

- Domínio 1: (6 itens) 4, 5, 7, 9, 12, 21
- Domínio 2: (4 itens) 6, 8, 16, 20
- Domínio 3: (5 itens) 3, 18, 23, 26, 30
- Domínio 4: (4 itens) 11, 22, 24, 29
- Domínio 5: (6 itens) 1, 15, 17, 19, 25, 27
- Domínio 6: (5 itens) 2, 10, 13, 14, 28

#### Interpretação dos escores:

- Domínio 1: 0-9 baixo nível de estresse; 10-12 médio nível de estresse; 13-14 alto nível de estresse; 15-18 muito alto nível de estresse.
- Domínio 2: 0-5 baixo nível de estresse; 6 médio nível de estresse; 7-8 alto nível de estresse; 9-12 muito alto nível de estresse.
- Domínio 3: 0-10 baixo nível de estresse; 11-12 médio nível de estresse; 13-14 alto nível de estresse; 15 muito alto nível de estresse.
- Domínio 4: 0-7 baixo nível de estresse; 8-10 médio nível de estresse; 11 alto nível de estresse; 12 muito alto nível de estresse.

estresse.

Domínio 5: 0-9 baixo nível de estresse; 10 médio nível de estresse; 11-12 alto nível de estresse; 13-18 muito alto nível de estresse.

Domínio 6: 0-9 baixo nível de estresse; 10-11 médio nível de estresse; 12-13 alto nível de estresse; 14-15 muito alto nível de estresse.

**Annex 3 - Stress Assessment in Nursing Students – Version Reduced [14]apud[15].**

Leia atentamente cada item abaixo e marque com um “X” o número correspondente com a intensidade de estresse que a situação lhe provoca, conforme a seguir:

0	1	2	3
<b>Não vivencio a situação</b>	<b>Não me sinto estressado com a situação</b>	<b>Me sinto pouco estressado com a situação</b>	<b>Me sinto muito estressado com a situação</b>

1	A obrigatoriedade em realizar os trabalhos extraclasse	0	1	2	3
2	Realizar os procedimentos assistenciais de modo geral	0	1	2	3
3	Comunicação com os demais profissionais da unidade de estágio	0	1	2	3
4	O ambiente da unidade clínica de estágio	0	1	2	3
5	Comunicação com os profissionais de outros setores no local de estágio	0	1	2	3
6	Ter medo de cometer erros durante a assistência ao paciente	0	1	2	3
7	A forma adotada para avaliar o conteúdo teórico	0	1	2	3
8	Distância entre a faculdade e o local de moradia	0	1	2	3
9	Sentir insegurança ou medo ao fazer as provas teóricas	0	1	2	3
10	O grau de dificuldade para a execução dos trabalhos extraclasse	0	1	2	3
11	Perceber as dificuldades que envolvem o relacionamento com outros profissionais da área	0	1	2	3
12	Perceber a responsabilidade profissional quando está atuando no campo de estágio	0	1	2	3
13	Observar atitudes conflitantes em outros profissionais	0	1	2	3
14	Transporte público utilizado para chegar à faculdade	0	1	2	3
15	Tempo exigido pelo professor para a entrega das atividades extraclasse	0	1	2	3
16	Distância entre a maioria dos campos de estágio e o local de moradia	0	1	2	3
17	Faltar tempo para o lazer	0	1	2	3
18	Transporte público utilizado para chegar ao local do estágio	0	1	2	3
19	Faltar tempo para momentos de descanso	0	1	2	3

**Figures 1 e 2 - Perceived Stress Scale (PSS) [16].**

Figure 1

**APPENDIX A:**  
**Items and Instructions for Perceived Stress Scale**

---

The questions in this scale ask you about your feelings and thoughts during the last month. In each case, you will be asked to indicate *how often* you felt or thought a certain way. Although some of the questions are similar, there are differences between them and you should treat each one as a separate question. The best approach is to answer each question fairly quickly. That is, don't try to count up the number of times you felt a particular way, but rather indicate the alternative that seems like a reasonable estimate.

For each question choose from the following alternatives:

- 0. never
- 1. almost never
- 2. sometimes
- 3. fairly often
- 4. very often

1. In the last month, how often have you been upset because of something that happened unexpectedly?
2. In the last month, how often have you felt that you were unable to control the important things in your life?
3. In the last month, how often have you felt nervous and "stressed"?
- 4.<sup>a</sup> In the last month, how often have you dealt successfully with irritating life hassles?
- 5.<sup>a</sup> In the last month, how often have you felt that you were effectively coping with important changes that were occurring in your life?
- 6.<sup>a</sup> In the last month, how often have you felt confident about your ability to handle your personal problems?
- 7.<sup>a</sup> In the last month, how often have you felt that things were going your way?
8. In the last month, how often have you found that you could not cope with all the things that you had to do?
- 9.<sup>a</sup> In the last month, how often have you been able to control irritations in your life?
- 10.<sup>a</sup> In the last month, how often have you felt that you were on top of things?

Figure 2

**APPENDIX A (Continued)**

---

11. In the last month, how often have you been angered because of things that happened that were outside of your control?
12. In the last month, how often have you found yourself thinking about things that you have to accomplish?
- 13.<sup>a</sup> In the last month, how often have you been able to control the way you spend your time?
14. In the last month, how often have you felt difficulties were piling up so high that you could not overcome them?

<sup>a</sup> Scored in the reverse direction.

Annex4 -Perceived Stress Scale in Brazilian Portuguese [16]apud[17].

**Itens e instruções para aplicação**

As questões nesta escala perguntam sobre seus sentimentos e pensamentos durante o último mês. Em cada caso, será pedido para você indicar o quanto frequentemente você tem se sentido de uma determinada maneira. Embora algumas das perguntas sejam similares, há diferenças entre elas e você deve analisar cada uma como uma pergunta separada. A melhor abordagem é responder a cada pergunta razoavelmente rápido. Isto é, não tente contar o número de vezes que você se sentiu de uma maneira particular, mas indique a alternativa que lhe pareça como uma estimativa razoável. Para cada pergunta, escolha as seguintes alternativas:

- 0= nunca
- 1= quase nunca
- 2= às vezes
- 3= quase sempre

4= sempre

Neste último mês, com que frequência...						
		0	1	2	3	4
1	Você tem ficado triste por causa de algo que aconteceu inesperadamente?	0	1	2	3	4
2	Você tem se sentido incapaz de controlar as coisas importantes em sua vida?	0	1	2	3	4
3	Você tem se sentido nervoso e “estressado”?	0	1	2	3	4
4	Você tem tratado com sucesso dos problemas difíceis da vida?	0	1	2	3	4
5	Você tem sentido que está lidando bem as mudanças importantes que estão ocorrendo em sua vida?	0	1	2	3	4
6	Você tem se sentido confiante na sua habilidade de resolver problemas pessoais?	0	1	2	3	4
7	Você tem sentido que as coisas estão acontecendo de acordo com a sua vontade?	0	1	2	3	4
8	Você tem achado que não consegue lidar com todas as coisas que você tem que fazer?	0	1	2	3	4
9	Você tem conseguido controlar as irritações em sua vida?	0	1	2	3	4
10	Você tem sentido que as coisas estão sob o seu controle?	0	1	2	3	4
11	Você tem ficado irritado porque as coisas que acontecem estão fora do seu controle?	0	1	2	3	4
12	Você tem se encontrado pensando sobre as coisas que deve fazer?	0	1	2	3	4
13	Você tem conseguido controlar a maneira como gasta seu tempo?	0	1	2	3	4
14	Você tem sentido que as dificuldades se acumulam a ponto de você acreditar que não pode superá-las?	0	1	2	3	4

#### Annex 5 -Perceived Stress Scale in Portuguese from Portugal [16]apud [18]

##### Anexo

##### Instruções e itens traduzidos da Perceived Stress Scale

Para cada questão, pedimos que indique com que frequência se sentiu ou pensou de determinada maneira, durante o último mês. Apesar de algumas perguntas serem parecidas, existem diferenças entre elas e deve responder a cada uma como perguntas separadas. Responda de forma rápida e espontânea. Para cada questão, escolha a alternativa que melhor se ajusta à sua situação.

**0 - Nunca. 1 - Quase nunca. 2 - Algumas vezes. 3 - Frequentemente. 4 - Muito frequentemente.**

1. No último mês, com que frequência esteve preocupado(a) por causa de alguma coisa que aconteceu inesperadamente?
2. No último mês, com que frequência se sentiu incapaz de controlar as coisas importantes da sua vida?
3. No último mês, com que frequência se sentiu nervoso(a) e em estresse?
- 4†. No último mês, com que frequência sentiu confiança na sua capacidade para enfrentar os seus problemas pessoais?
- 5†. No último mês, com que frequência sentiu que as coisas estavam acorrer à sua maneira?
6. No último mês, com que frequência sentiu que não aguentava com as coisas todas que tinha para fazer?
- 7†. No último mês, com que frequência foi capaz de controlar as suas irritações?
- 8†. No último mês, com que frequência sentiu ter tudo sob controlo?
9. No último mês, com que frequência se sentiu furioso(a) por coisas que ultrapassaram o seu controlo?
10. No último mês, com que frequência sentiu que as dificuldades se estavam a acumular tanto que não as conseguia ultrapassar?

† Itens cotados de forma inversa.

#### Annex 6 - Perceived Stress Scale 4 (PSS-4)[16] apud [19].

#### Perceived Stress Scale 4(PSS-4)

#### INSTRUCTIONS

The questions in this scale ask you about your feelings and thoughts during THE LAST MONTH. In each case, please indicate your response by placing an "X" over the square representing HOW OFTEN you felt or thought a certain way.

	Never	Almost Never	Sometimes	Fairly Often	Very Often
	0	1	2	3	4
1. In the last month, how often have you felt that you were unable to control the important things in your life?	<input type="checkbox"/>				
2. In the last month, how often have you felt that you were unable to control the important things in your life?	<input type="checkbox"/>				
3. In the last month, how often have you felt confident about your ability to handle your personal problems?	<input type="checkbox"/>				
4. In the last month, how often have you felt that things were going your way?	<input type="checkbox"/>				
5. In the last month, how often have you felt difficulties were piling up so high that you could not overcome them?	<input type="checkbox"/>				

#### Scoring for the Perceived Stress Scale 4:

##### Questions 1 and 4

0 = Never

1 = Almost Never

2 = Sometimes

3 = Fairly Often

4 = Very Often

##### Questions 2 and 3

4 = Never

3 = Almost Never

2 = Sometimes

1 = Fairly Often

0 = Very Often

Lowest score: 0

Highest score: 16

Higher scores are correlated to more stress.

**Annex7** -Lipp Adult Stress Symptom Inventory [20] *apud* [1].

#### QUADRO - 1

#### QUADRO - 2

Você vai me dizer os sintomas que sentiu na última semana (de \_\_\_\_\_ até hoje)

- ( ) 01 Problemas com amemória
- ( ) 02 Mal-estar generalizado, sem motivo
- ( ) 03 Formigamento demãos/pés
- ( ) 04 Sensação de desgaste físicoconstante
- ( ) 05 Mudança de appetite
- ( ) 06 Aparecimento de problema de pele
- ( ) 07 Pressão alta
- ( ) 08 Cansaço constante
- ( ) 09 Aparecimento de úlcera
- ( ) 10 Tontura ou sensação de estar flutuando
- ( ) 11 Muito sensível em nível deemoção
- ( ) 12 Dúvida quanto a si próprio
- ( ) 13 Pnesar direto em um só assunto
- ( ) 14 Irritabilidade excessiva

### QUADRO 3

Você vai me dizer os sintomas que sentiu no último mês (deaté hoje).

- ( ) 01 Diarreia frequente
- ( ) 02 Dificuldade com sexo
- ( ) 03 Dificuldade para pegar no sono/acordar durante a noite
- ( ) 04 Náuseas/ânsia de vômito
- ( ) 05 Tiques/manias, por exemplo: ficar mexendo no cabelo
- ( ) 06 Pressão alta direto
- ( ) 07 Problemas de pele por um tempo longo
- ( ) 08 Mudança extrema de apetite
- ( ) 09 Excesso de gases [estômago/intestino(barriga)]
- ( ) 10 Tontura frequente
- ( ) 11 Úlcera
- ( ) 12 Enfarte
- ( ) 13 Impossibilidade de trabalhar ( ) 14 Pesadelos
- ( ) 15 Sensação de não ser competente em todas as áreas

#### Avaliação

F1 ( )  
F2 ( )  
F3 ( )  
Total (vertical)

**Total (horizontal)**  
P1 ( )  
P2 ( )  
P3 ( )  
P( )