

Age and Nutritional Status by Mini-Nutritional Assessment (MNA) can discriminate Alzheimer's Disease Severity

Vania Aparecida Leandro-Merhi^{1,*}, Tamires Barbosa Nascimento dos Santos², Gloria Maria de Almeida Souza Tedrus³

¹PhD, Professor Doctor of the Postgraduate Program in Health Sciences, PUC-Campinas-SP-Brazil.

²Ms, Postgraduate Program in Health Science, Puc-Campinas-SP-Brazil.

³PhD, Professor Doctor of the Postgraduate Program in Health Sciences, PUC-Campinas-SP-Brazil.

*Corresponding author

Received: 06 Dec 2020;

Received in revised form:

03 Feb 2021;

Accepted: 27 Feb 2021;

Available online: 28 Mar 2021

©2021 The Author(s). Published by AI Publication. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords— *Alzheimer disease, anthropometry, mini-nutritional assessment, associated factors.*

Abstract— *Investigate the risk factors associated with Clinical Dementia Rating (CDR) in patients with Alzheimer's disease (AD). Cross-sectional study with 43 elderly people with AD. Parameters assessing nutritional status (anthropometry and mini-nutritional assessment-MNA), neurocognitive assessment and Clinical Dementia Rating (CDR) were investigated. For the investigation of AD severity risk factors, statistical analysis was performed using logistic regression models. The level of significance adopted was 5%. No statistically significant difference in the relationship between the parameters of anthropometry nutritional assessment and the CDR score were found. In the final regression model; only MNA ($p=0.0159$; $OR=4.815$; $IC95\%=1.342$; 17.282) and age ($p=0.0481$; $OR= 1.097$; $IC95\%=1.001$; 1.202), were associated with the severity of the disease (CDR). Age and MNA discriminate the severity of the disease and were considered risk factors for the severity of AD.*

I. INTRODUCTION

Alzheimer's disease (AD) is a neurodegenerative disease prevalent in the elderly. It is well known that incorporation of lifestyle changes, as well as physical and recreational activities, diet and cognitive activity, among others, are important to prevent the disease¹. Evidence of preventive interventions¹ could encourage or contribute to the treatment of this population. Recent studies^{2,3} have indicated the importance of some nutrients which shortage is associated with the development of AD. These nutrients contribute to the synthesis of neurotransmitters, modulation in epigenetic mechanisms and with antioxidants². In elderly patients with subjective cognitive decline, clinically relevant findings have been associated with the quality of the diet³.

Regarding the nutritional status of patients with AD, a study⁴ showed a high prevalence of malnutrition or risk of malnutrition detected by the mini nutritional assessment (MAN), in patients with different severity, also associated with reduced functional status. Another recent study⁵ which investigated the association between nutritional status and clinical and cognitive aspects of patients with AD, showed significant progressive impairment of nutritional and cognitive indicators related to disease progression. In a cross-sectional study⁶, assessing the nutritional status and body composition and mild cognitive impairment in patients with AD, the MAN score was significantly lower, compared to healthy elderly controls. Dementia was also considered an independent predictor of malnutrition risk, when compared to cognitively intact individuals⁷.

Recently, a prospective multicenter study⁸ investigating the nutritional status in AD and its influence on disease progression, found that nutritional impairment was highly prevalent in patients with AD. The authors also pointed out that the identification of the nutritional status in early diagnosis, could help those patients who are at a higher risk of disease progression.⁸ The relevance of assessing the nutritional status with different indicators in patients with AD, was also reported in other investigations^{9,10,11}.

In view of all the findings discussed above, this study investigated nutritional risk factors using the clinical dementia rating (CDR), in patients with Alzheimer's disease (AD).

II. CASES AND METHOD

This study is part of a more comprehensive research project on lifestyle aspects in elderly patients with dementia. Some nutritional aspects have been addressed in another paper⁵.

This was a cross-sectional study, which included outpatient elderly individuals (n=43) aged ≥ 65 years with diagnosis of suspected DA, according to the Diagnostic and Statistical Manual of Mental Disorders¹², the recommendations of the European Federation of Neurological Societies¹³ and the Brazilian Academy of Neurology¹⁴. Clinical dementia rating (CDR)¹⁵ was used to stage the severity of dementia, that was classified as mild, moderate or severe. Patients with other serious illnesses and patients and/or caregivers unable to answer the evaluation questions were excluded. The study was approved by the institution's ethics research committee (No. 1,234,677) and was started after the guardians signed the free and informed consent form.

Methodological procedures:

a) *Nutritional assessment by anthropometry*: According to standardized procedures and cutoff points reported in the literature such as the body mass index (BMI) for the elderly¹⁶, the calf circumference (CC)¹⁷, the thickness of the adductor pollicis muscle (APM)^{18,19}, the handgrip strength (HGS)^{20,21,22} and waist circumference^{23,24}. The anthropometric indicators of body composition^{17,25} of arm

circumference (AC), triceps skinfold (TSF), arm muscle circumference (AMC) and sub-scapular skinfold (SSF), were classified in percentile distribution as: lower than the 5th percentile (<P5: depletion); between the 5th percentile and below the 10th percentile (P5 - <P10: risk of depletion); between 10th and 90th percentiles (P10 - P90: normal weight); above the 90th percentile and up to the 95th percentile (> P90 - P95: risk for excess) and above the 95th percentile (> P95: excess).

b) *Mini Nutritional Assessment (MNA)*: The MNA questionnaire was applied and the nutritional status was classified according to the cutoff points established as eutrophic, risk of malnutrition and malnutrition²⁶.

c) *International physical activity questionnaire - short form (IPAQ-SF)*: Physical activity was assessed according to a standardized and validated instrument for the Brazilian population²⁷. For the purpose of analysis, in this study, physical activity was classified as: sedentary or active.

d) *Statistical analysis*: A descriptive analysis was performed with frequency tables for categorical variables and measures of position and dispersion for continuous variables. The Chi-square or Fisher's exact test was used, when necessary; for comparing proportions. To compare continuous or orderable measurements between 2 groups, the Mann-Whitney test was applied and among 3 groups, the Kruskal-Wallis test. Later, to identify the predictors of disease severity by the CDR clinical score, regression models were used, such as univariate and multiple logistic regression analysis. The variable selection process employed was stepwise. The level of significance adopted for the statistical tests was 5%.

III. RESULTS

In our study's population (n=43), 16 (37.2%) patients were rated by CDR in the mild stage; 19 (44.1%) patients in the moderate stage and 8 (18.6%) patients in the severe stage. The sample included 28 (65.1%) female patients and 15 (34.9%) male patients.

There was no statistically significant difference in the relationship between all anthropometric parameters, age and IPAQ-SF with the CDR scores (Table 1).

Table 1. Descriptive analysis of the studied variables and comparisons with the CDR score.

Variables	Classification		CDR - L N=16	CDR - M N=19	CDR - G N=8	Total N=43	P-value
Age	years	x±dp	77.6 ± 6.0	82.2 ± 7.5	82.8 ± 6.3	80.6 ± 7.0	0.0959 ¹
Weight	Kg	x±dp	65.8 ± 13.8	61.5 ± 13.5	55.2 ± 9.2	62.0 ± 13.2	0.1728 ¹

Height	cm	x±dp	155.9 ± 10.3	152.4 ± 8.8	154.6 ± 9.4	154.1 ± 9.4	0.6743 ¹
BMI	kg/m ²	x±dp	26.9 ± 4.1	26.3 ± 4.5	23.2 ± 3.9	26.0 ± 4.4	0.1129 ¹
HGS-bd	Kg	x±dp	16.5 ± 10.1	12.2 ± 7.2	12.8 ± 8.2	13.9 ± 8.6	0.4745 ¹
HGS-be	Kg	x±dp	15.6 ± 9.1	11.4 ± 7.2	11.4 ± 7.1	13.0 ± 8.0	0.3847 ¹
HGS average	Kg	x±dp	16.0 ± 9.6	11.8 ± 7.1	12.1 ± 7.6	13.4 ± 8.2	0.4618 ¹
AC	P10 – P90	n (%)	7 (43.8)	8 (42.1)	4 (50.0)	19 (44.2)	1.0000 ³
	>P90 – P95	n (%)	9 (56.3)	11 (57.9)	4 (50.0)	24 (55.8)	
TSF	P10 – P90	n (%)	8 (50.0)	10 (52.6)	4 (50.0)	22 (51.2)	0.9854 ²
	>P90 – P95	n (%)	8 (50.0)	9 (47.4)	4 (50.0)	21 (48.8)	
AMC	P10 – P90	n (%)	9 (56.3)	11 (57.9)	8 (100.0)	28 (65.1)	0.0715 ²
	>P90 – P95	n (%)	7 (43.8)	8 (42.1)	0 (0.0)	15 (34.9)	
SSF	P5 – <P10	n (%)	1 (6.3)	2 (10.5)	0 (0.0)	3 (7.0)	1.0000 ³
	P10 – P90	n (%)	15 (93.8)	17 (89.5)	8 (100.0)	40 (93.0)	
CC (cm)	>31 cm	n (%)	13 (81.3)	14 (73.7)	4 (50.0)	31 (72.1)	0.3240 ³
	≤31 cm	n (%)	3 (18.8)	5 (26.3)	4 (50.0)	12 (27.9)	
WC (cm)	no risk	n (%)	2 (12.5)	4 (21.1)	1 (12.5)	7 (16.3)	0.8653 ³
	with risk	n (%)	14 (87.5)	15 (78.9)	7 (87.5)	36 (83.7)	
IMC (kg/m ²)	22 – 27	n (%)	7 (43.8)	9 (47.4)	5 (62.5)	21 (48.8)	0.6600 ³
	<22	n (%)	2 (12.5)	3 (15.8)	2 (25.0)	7 (16.3)	
	>27	n (%)	7 (43.8)	7 (36.8)	1 (12.5)	15 (34.9)	

MNA	eutrophic	n (%)	10 (62.5)	8 (42.1)	1 (12.5)	19 (44.2)	0.0658 ³
	MR	n (%)	6 (37.5)	11 (57.9)	7 (87.5)	24 (55.8)	
TAPM (mm)	minimum (7 or 5)	n (%)	15 (93.8)	18 (100.0)	7 (87.5)	40 (95.2)	-
	median (12 or 11)	n (%)	1 (6.3)	0 (0.0)	1 (12.5)	2 (4.8)	
	maximum (20 or 18)	-	-	-	-	-	
CLAPA	sedentary	n (%)	9 (56.3)	11 (57.9)	6 (75.0)	26 (60.5)	0.7758 ³
	active	n (%)	7 (43.8)	8 (42.1)	2 (25.0)	17 (39.5)	

¹ Kruskal-Wallis test; ² Chi-square test; ³ Fisher's exact test.

Legend:- CDR: clinical dementia rating; BMI: body mass index; HGSra: hand grip strength - right arm; HGSla: hand grip strength-left arm; HGS Max: maximum handgrip strength; AC: arm circumference; TSF: tricipital skin fold; AMC: arm muscle circumference; SSF: subscapular skin fold; CC: calf circumference; WC: waist circumference; MNA: mini nutritional assessment; MR: risk of malnutrition, TAPM: thickness of the adductor pollicis muscle; CLAPA: classified physical activity. P: percentile.

CDR - L: clinical dementia rating - mild; CDR-M: clinical dementia rating - moderate; CDR - G: clinical dementia rating - serious

In the analysis among all the anthropometry variables studied (Table 2). The other anthropometric data parameters and physical activity classified by IPAQ, no significant statistical difference was found in all the variables studied (Table 2). The other anthropometric data were not significantly related to the practice of Physical Activities in AD elderly (Table 2).

Table 2. Descriptive analysis of the variables studied and comparisons with physical activity by the IPAQ classified.

Variables	Classification		PA sedentary N=26	PA active N=17	Total N=43	P-value
Age	years	x±dp	80.1 ± 7.6	81.3 ± 6.2	80.6 ± 7.0	0.7843 ¹
Weight	Kg	x±dp	61.1 ± 13.7	63.3 ± 12.7	62.0 ± 13.2	0.5594 ¹
Height	cm	x±dp	153.3 ± 10.5	155.2 ± 7.4	154.1 ± 9.4	0.4486 ¹
BMI	kg/m ²	x±dp	25.9 ± 4.7	26.1 ± 4.0	26.0 ± 4.4	0.8913 ¹
HGS-bd	Kg	x±dp	14.0 ± 8.5	13.8 ± 9.0	13.9 ± 8.6	0.9702 ¹
HGS-be	Kg	x±dp	13.0 ± 7.4	12.9 ± 9.0	13.0 ± 8.0	0.8423 ¹
HGS average	Kg	x±dp	13.5 ± 7.9	13.4 ± 9.0	13.4 ± 8.2	0.8815 ¹

AC	P10 – P90	n (%)	10 (38.5)	9 (52.9)	19 (44.2)	0.3499 ²
	>P90 – P95	n (%)	16 (61.5)	8 (47.1)	24 (55.8)	
TSF	P10 – P90	n (%)	12 (46.2)	10 (58.8)	22 (51.2)	0.4164 ²
	>P90 – P95	n (%)	14 (53.8)	7 (41.2)	21 (48.8)	
AMC	P10 – P90	n (%)	16 (61.5)	12 (70.6)	28 (65.1)	0.5427 ²
	>P90 – P95	n (%)	10 (38.5)	5 (29.4)	15 (34.9)	
SSF	P5 – <P10	n (%)	1 (3.8)	2 (11.8)	3 (7.0)	0.5523 ³
	P10 – P90	n (%)	25 (96.2)	15 (88.2)	40 (93.0)	
CC (cm)	>31 cm	n (%)	17 (65.4)	14 (82.4)	31 (72.1)	0.3065 ³
	≤31 cm	n (%)	9 (34.6)	3 (17.6)	12 (27.9)	
WC (cm)	no risk	n (%)	4 (15.4)	3 (17.6)	7 (16.3)	1.0000 ³
	with risk	n (%)	22 (84.6)	14 (82.4)	36 (83.7)	
IMC (kg/m ²)	22 – 27	n (%)	12 (46.2)	9 (52.9)	21 (48.8)	0.9154 ³
	<22	n (%)	4 (15.4)	3 (17.6)	7 (16.3)	
	>27	n (%)	10 (38.5)	5 (29.4)	15 (34.9)	
MNA	euthrophic	n (%)	9 (34.6)	10 (58.8)	19 (44.2)	0.1181 ²
	MR	n (%)	17 (65.4)	7 (41.2)	24 (55.8)	

¹ Mann-Whitney test, ² Chi-square test, ³ Fisher's exact test

Legend:- PA: physical activity, BMI: body mass index; HGS-ra: hand grip strength-right arm; HGS-la: hand grip strength-left arm; HGS: handgrip strength; AC: arm circumference; TSF: tricipital skin fold; AMC: muscle circumference of the arm; SSF: subscapular skin fold; CC: calf circumference; WC: waist circumference; P: percentile; MNA: mini nutritional assessment; MR: malnutrition risk.

When investigating the risk factors associated with the CDR score, using univariate and multiple logistic regression analysis, it was found that in the final model, age (p=0.0481; OR=1.097; 95% CI=1.001; 1,202) and the assessment of nutritional status by MNA (p=0.0159; OR=4,815; 95% CI=1,342; 17,282), were the factors that, together, better discriminated the

severity of the disease. Each additional year of age increased by 9.7% the chance of CDR score 2 or 3. The patients' nutritional status by MNA, yielded patients classified as malnourished or at risk of malnutrition with a chance of being a score 2 or 3, 4.8 times higher when compared to the nutritional status of normal weight (Table 3).

Table 3. Risk factors associated with the CDR score, investigated by univariate and multiple logistic regression analysis.

<i>Univariate Analysis</i>				
Variable	Reference	P-value	OR	CI (95%)
Age		0.0564	1.088	0.998; 1.187
BMI		0.0896	0.888	0.775; 1.018
HGS		0.1504	0.948	0.882; 1.019
AC		0.1355	0.897	0.779; 1.034
AMC		0.3214	0.990	0.970; 1.010
SSF		0.2401	0.946	0.862; 1.038
WC		0.4333	0.980	0.932; 1.031
CC		0.1137	0.870	0.732; 1.034
TAPM		0.4622	0.926	0.754; 1.137
TSF	eutrophy versus ER + excess	0.9576	1.031	0.336; 3.158
SSF	Depletion + DPR versus eutrophy + excess	0.8051	0.757	0.083; 6.920
AC	eutrophy versus ER +excess	0.8405	1.123	0.364; 3.466
CC	eutrophy versus malnutrition	0.1344	0.375	0.104; 1.354
AMC	eutrophy versus ER +excess	0.0976	2.817	0.827; 9.593
IMC	eutrophy versus overweight	0.2427	2.132	0.599; 7.592
	thinness versus overweight	0.2469	2.747	0.497; 15.204
WC	no risk versus with risk	0.8261	1.185	0.261; 5.386
MNA	MR+D	0.0248	4.006	1.192; 13.464
	MA versus eutrophy			
CLAPA	sedentary versus active			
<i>Multiple analysis*</i>				
Variable	Reference	P-value	OR	CI (95%)
Age		0.0481	1.097	1.001; 1.202
MNA	MR+MA versus eutrophy	0.0159	4.815	1.342; 17.282

Legend:- CDR: clinical dementia rating; BMI: body mass index; HGS: handgrip strength; AC: arm circumference; AMC: arm muscle circumference; SSF: subscapular skin fold; DPR: depletion risk; ER: excess risk; WC: waist circumference; CC: calf circumference; TAPM: thickness of the adductor pollicis muscle; TSF: triceps skin fold; MNA: mini nutritional assessment; MAR malnutrition risk; MA: malnourished; CLAPA: classified physical activity.

* multiple model: stepwise process, model accuracy (statistics): c = 0.725.

IV. DISCUSSION

This investigation used several anthropometric parameters to assess the nutritional status of elderly people

with AD, and the initial hypothesis of this investigation was to find an association between the different anthropometric parameters used and the disease severity

score (CDR), yet this association was not observed. In the regression models, investigating what the risk factors associated with the severity of the disease would be, only age and MNA remained in the final model. The MNA addresses in its assessment instrument, the indicators of arm and calf circumference, functional capacity, dietary and body weight changes, nutritional problems, disease, elderly people physical evaluation ²⁶. And, the present study pointed out MNA (risk of malnutrition and malnutrition), as a predictor of the severity of the disease in elderly people with AD. The pathophysiological mechanisms between nutritional status and AD are not yet fully understood. Our data suggest that older age and risk of malnutrition and malnutrition and nutritional imbalance conditions are associated with greater severity of dementia in elderly people with AD. An association between nutritional status and the rate of cognitive and functional decline was also observed in an observational longitudinal study ¹¹, using MNA, CDR and the Mini-Mental State Exam. In the study in question, it was found that a worsening of the CDR, during dementia, was associated with lower MNA scores ¹¹.

Other studies suggest an association between a greater impairment of nutritional status with more advanced AD stages, faster cognitive and functional decline and higher mortality rates ^{4,8,28}.

Unlike the findings found in this investigation, a prospective study recently carried out in Portugal ²⁹ with elderly patients with AD, showed that malnutrition, unintentional weight loss, low weight, low values of calf circumference and muscle circumference of the arm, of handgrip strength and gait speed, were associated with a higher risk of death, regardless of gender, age, marital status, education and cognitive function status²⁹. In our study, among the investigated nutritional variables, only malnutrition was associated with AD. And in a Korean study ⁷ that investigated the nutritional status and clinical risk factors of malnourished patients, the authors observed that dementia was considered an independent predictor for the risk of malnutrition, further suggesting that cortical thinning in the left temporal regions, could be related to nutritional status. Another recent study with female patients with mild cognitive impairment and early-stage Alzheimer's disease, found malnutrition associated with behavioral and psychiatric symptoms of dementia ³⁰.

Other recent findings ³¹ suggest that malnutrition may be related not only to impaired cognition, but also to the pathology of AD. The authors pointed out that AD biomarkers were associated with the MNA score, waist circumference and BMI, and the

associations with the MNA score remained after adjustment for cognitive performance ³¹.

The present study found no association between the severity of the disease and the HGS. Different data were recently reported in a study with an elderly Japanese population, which showed a greater decline in HGS in the elderly, associated with the late onset of dementia ²¹.

As limiting factors of this study, it is highlighted that our service is a university hospital, but not a tertiary public health center in Brazil. This study is cross-sectional, with a relatively small sample of cases, although it is a specific population with elderly population with AD, monitored on an outpatient basis. However, we must consider that the hospital and outpatient service, where our study was conducted, is part of a representative clinical and surgical neurology service that includes AD patients care in a large metropolitan region. Thus, the individuals who participated in this study were representative of all those who undergo routine clinical treatment, besides the fact that this population underwent specific screening that allowed diagnostic tracking of the severity of the disease. These points can be considered the strong points of our investigation. Anyway, further investigations with a larger sample size are still need to assess the impact of these findings. We can also suggest that nutritional counseling be offered to those patients who are at risk of malnutrition.

V. CONCLUSION

The findings in this study allowed us to conclude that age and MNA, discriminate the severity of AD and were considered risk factors for the severity of the disease.

REFERENCES

- [1] Schelke MW, Hackett K, Chen JL, Shih C, Shum J, Montgomery ME, Chiang GC, Berkowitz C, Seifan A, Krikorian R, Isaacson RS. Nutritional interventions for Alzheimer's prevention: a clinical precision medicine approach. *Ann N Y Acad Sci.* 2016 Mar; 1367(1):50-6. doi: 10.1111/nyas.13070.
- [2] Muñoz Fernández SS, Lima Ribeiro SM. Nutrition and Alzheimer Disease. *Clin Geriatr Med.* 2018 Nov; 34(4):677-697. doi: 10.1016/j.cger.2018.06.012.
- [3] Wesselman LMP, Doorduijn AS, de Leeuw FA, Verfaillie SCJ, van Leeuwenstijn-Koopman M, Slot RER, Kester MI, Prins ND, van de Rest O, de van der Schueren MAE, Scheltens P, Sikkes SAM, van der Flier WM. Dietary Patterns Are Related to Clinical Characteristics in Memory Clinic Patients with Subjective Cognitive Decline: The SCIENCE Project. *Nutrients.* 2019 May 11;11(5). pii: E1057. doi: 10.3390/nu11051057.

- [4] Tombini M, Sicari M, Pellegrino G, Ursini F, Insardá P, Di Lazzaro V. Nutritional Status of Patients with Alzheimer's Disease and Their Caregivers. *J Alzheimers Dis.* 2016 Oct 18;54(4):1619-1627.
- [5] Santos TBND, Fonseca LC, Tedrus GMAS, Delbue JL. Alzheimer's disease: nutritional status and cognitive aspects associated with disease severity. *Nutr Hosp.* 2018 Dec 3;35(6):1298-1304. doi: 10.20960/nh.2067.
- [6] Cova I, Pomati S, Maggiore L, Forcella M, Cucumo V, Ghiretti R, Grande G, Muzio F, Mariani C. Nutritional status and body composition by bioelectrical impedance vector analysis: A cross sectional study in mild cognitive impairment and Alzheimer's disease. *PLoS One* 2017 Feb 10;12(2):e0171331. doi: 10.1371/journal.pone.0171331. eCollection 2017.
- [7] Jang JW, Kim Y, Choi YH, Lee JM, Yoon B, Park KW, Kim SE, Kim HJ, Yoon SJ, Jeong JH, Kim EJ, Jung NY, Hwang J, Kang JH, Hong JY, Choi SH. Association of Nutritional Status with Cognitive Stage in the Elderly Korean Population: The Korean Brain Aging Study for the Early Diagnosis and Prediction of Alzheimer's Disease. *J Clin Neurol.* 2019 Jul;15(3):292-300. doi: 10.3988/jcn.2019.15.3.292.
- [8] Izquierdo Delgado E, Gutiérrez Ríos R, Andrés Calvo M, Repiso Gento I, Castrillo Sanz A, Rodríguez Herrero R, Rodríguez Sanz MF, Tola-Arribas MA. Nutritional status assessment in Alzheimer disease and its influence on disease progression. *Neurologia* 2020 Jan 21. pii: S0213-4853(19)30148-3. doi: 10.1016/j.nrl.2019.11.005.
- [9] Ivanski F, Nascimento LP, Fermio BL, Bonini JS, Silva WCFND, Valério JMS, Fabbri R, Bosetto AK, Gregório E. Nutritional evaluation of geriatric patients with Alzheimer's disease in Southern Brazil: case-control study. *Nutr Hosp.* 2018 Apr 5;35(3):564-569. doi: 10.20960/nh.1626.
- [10] Sun J, Wen S, Zhou J, Ding S. Association between malnutrition and hyperhomocysteine in Alzheimer's disease patients and diet intervention of betaine. *J Clin Lab Anal.* 2017 Sep; 31(5). doi: 10.1002/jcla.22090.
- [11] Sanders C, Behrens S, Schwartz S, Wengreen H, Corcoran CD, Lyketsos CG, Tschanz JT. Nutritional Status is Associated with Faster Cognitive Decline and Worse Functional Impairment in the Progression of Dementia: The Cache County Dementia Progression Study 1. *J Alzheimers Dis.* 2016 Feb 27;52(1):33-42. doi: 10.3233/JAD-150528.
- [12] McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Kawas CH et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's Dement.* 2011;7(3):263-9.
- [13] Waldemar G, Dubois B, Emre M, Georges J, McKeith G, Rossor M et al. Recommendations for the diagnosis and management of Alzheimer's disease and other disorders associated with dementia: EFNS guideline. *Eur J Neurol.* 2007;14;14(1):1-26.
- [14] Nitrini R, Caramelli P, Bottino CMC, Damasceno BP, Brucki SMD, Anghinah R. Critérios diagnósticos e exames complementares. Recomendações do Departamento de Neurologia Cognitiva e do Envelhecimento da Academia Brasileira de Neurologia. *Arq Neuropsiquiatr.* 2005;63(3-A):713-9.
- [15] Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology* 1993;43:632-7.
- [16] Lipschitz DA. Screening for nutritional status in the elderly. *Prim Care* 1994; 22(1):55-67.
- [17] World Health Organization - WHO. Physical status: the use and interpretation of anthropometry. Report of a WHO expert consultation. Geneva: WHO Technical Report Series. 1995; 854:1-452.
- [18] Lameu EB, Gerude MF, Corrêa RC, Lima KA. Adductor pollicis muscle: a new anthropometric parameter. *Rev Hosp Clin Fac Med Sao Paulo* 2004; 59(2):57-62.
- [19] Bragagnolo R, Caporossi FS, Dock-Nascimento DB, Aguilar-Nascimento JE. Espessura do músculo adutor do polegar: um método rápido e confiável na avaliação nutricional de pacientes cirúrgicos. *Rev Col Bras Cir.* 2009; 36(5):371-6.
- [20] Dias JA, Ovando AC, Kulkamp W, Borges Junior NG. Força de prensão palmar: métodos de avaliação e fatores que influenciam a medida. *Rev Bras Cineantropom Desempenho Hum* 2010; 12(3):209-216.
- [21] Kishimoto H, Hata J, Ninomiya T, et al. Midlife and late life handgrip strength and risk of cause-specific death in a general Japanese population: the Hisayama Study. *J Epidemiol Community Health* 2014; 68:663-668.
- [22] Hatabe Y, Shibata M, Ohara T, et al. Decline in Handgrip Strength From Midlife to Late-Life is Associated With Dementia in a Japanese Community: The Hisayama Study. *J Epidemiol.* 2020; 30(1):15-23. doi:10.2188/jea.JE20180137
- [23] International Diabetes Federation – IDF. The IDF Consensus worldwide definition of the metabolic syndrome; 2005.
- [24] World Health Organization. Obesity: preventing and managing the global epidemic. Report of a WHO Consultation on obesity. Geneva: WHO, 1998.
- [25] Burr ML, Phillips MK. Anthropometric norms in the elderly. *British Journal of Nutrition* 1984; 51:165-9.
- [26] Guigoz Y, Garry JP. Mini nutritional assessment: A practical assessment tool for grading the nutritional state of elderly patients. *Facts and Research in Gerontology* 1994; Supplement (2):15-59.
- [27] Pardini R, Matsudo SM, Araujo T, Matsudo V, Andrade E, Braggion G, et al. Validação do questionário internacional de nível de atividade física (IPAQ - versão 6): Estudo piloto em adultos jovens brasileiros. *Rev Bras Cien Mov* 2001; 9:45-51.
- [28] Droogsma E, van Asselt DZ, Schölzel-Dorenbos CJ, van Steijn JH, van Walderveen PE, van der Hooft CS. Nutritional status of community-dwelling elderly with newly diagnosed Alzheimer's disease: prevalence of malnutrition and the relation of various factors to

nutritional status. *J Nutr Health Aging*. 2013; 17(7):606-610. doi:10.1007/s12603-013-0032-9

- [29] De Sousa OV, Mendes J, Amaral TF. Nutritional and Functional Indicators and Their Association With Mortality Among Older Adults With Alzheimer's Disease. *Am J Alzheimers Dis Other Demen*. 2020;35:1533317520907168. doi:10.1177/1533317520907168
- [30] Kimura A, Sugimoto T, Kitamori K, et al. Malnutrition is Associated with Behavioral and Psychiatric Symptoms of Dementia in Older Women with Mild Cognitive Impairment and Early-Stage Alzheimer's Disease. *Nutrients*. 2019;11(8):1951. Published 2019 Aug 20. doi:10.3390/nu11081951
- [31] Doorduijn AS, Visser M, van de Rest O, et al. Associations of AD Biomarkers and Cognitive Performance with Nutritional Status: The NUDAD Project. *Nutrients*. 2019;11(5):1161. Published 2019 May 23. doi:10.3390/nu11051161